

Review Article

Determinants of Child Undernutrition in Tanzania: Agriculture and Season Perspective: A Review

Happiness Samuel Muhimbula

Department of Human Nutrition and Consumer Sciences, College of Agriculture, Sokoine University of Agriculture, Morogoro, Tanzania

Email address:

happy.issa@sua.ac.tz

To cite this article:Happiness Samuel Muhimbula. (2024). Determinants of Child Undernutrition in Tanzania: Agriculture and Season Perspective: A Review. *Journal of Food and Nutrition Sciences*, 12(1), 1-19. <https://doi.org/10.11648/j.jfns.20241201.11>**Received:** November 7, 2023; **Accepted:** December 8, 2023; **Published:** January 8, 2024

Abstract: The causes of undernutrition are known to be complex and interrelated which need responses and solutions from different sectors to effectively design proper nutrition interventions. The UNICEF conceptual framework is a useful tool for understanding the causes of undernutrition and serves as a guide for assessing and analysing the causes, particularly among children in developing countries. Identifying determinants of child nutrition status is a necessary step in designing interventions, policies and programme that can support the scaling up of nutrition. This review scrutinizes undernutrition and its determinants based on the UNICEF conceptual framework on malnutrition. Poor feeding practices and characteristic of complementary foods were observed as immediate causes while household's food insecurity, environmental factors such as agricultural practices and seasonal influences as underlying determinants particularly in rural households. The highlight from this review suggests that seasons had significant impact on both immediate and underlying determinants of undernutrition and therefore intervention to improve nutrition status should consider variation in agriculture seasons. Lastly, multicomponent interventions focusing on agricultural interventions and practices that are affecting immediate changes in nutrition status among infants and young children are emphasized. Future attempt to improve children nutrition status in rural households should be multicomponent addressing nutrition sensitives farming practices including rural livestock intensification, poverty alleviation and considering agricultural seasons in all planning.

Keywords: Undernutrition, Feeding Practices, Nutrition-Sensitive Agriculture, Food Security, Seasons

1. Introduction

Child undernutrition remains a major global problem and is an underlying cause of 3.1 million child deaths annually [25, 160]. In an analysis accounting for several nutrition risks factors including; intrauterine growth restriction, stunting, wasting, deficiencies of vitamin A, Calcium, selenium, iron, folate and zinc and sub-optimum breastfeeding, 45% of all child deaths were attributed to undernutrition [25]. The root cause of most undernutrition and morbidity among infants and young children is poverty leading to an increasing burden of stunting and micronutrient deficiencies [13]. Undernutrition and micronutrient deficiencies often coexist, which is particularly detrimental to a child's life resulting in poor cognition, health and physical growth [12].

Child undernutrition results from a combination of

interrelated factors; however, the common denominator among all types of undernutrition is poor food and nutrition security, which includes poor infant feeding practices and poor dietary quality in early life [84]. In Tanzania, as in many developing countries, children consume and poor diet, mainly, cereal-based porridge with low nutrient densities [43, 52, 83]. Poor nutrition weakens children's immune systems, cognitive, and physiological growth, and is associated with increased child morbidity and mortality [38]. Climate variability, including seasonal food availability and poor rainfall distribution, increases the risk of morbidity and food insecurity, thereby affecting feeding practices and dietary patterns [13]. Additionally, poor maternal nutritional status during pregnancy and lactation has significant implications for poor nutrition outcome for children [21, 94]

The role of nutrition-sensitive agriculture in improving

nutrition particularly targeting the “first 1000 days”, has been a global focus Rural household are an ideal target for this type of intervention given the high rates of undernutrition and high dependency on agriculture for food consumption and livelihoods [101, 112]. Efforts to reduce undernutrition have been largely focused on nutrition specific interventions which reflect Western medical model, while agricultural, environmental and social factors which underpin the root cause of undernutrition has not been given much attention [28]. Although there is a strong relationship between agriculture and nutrition, there is a need to identify agricultural practices that can effect changes within the 1000 days when designing appropriate interventions. Tanzania is an agriculturally diverse country and there is substantial regional variation in the prevalence of undernutrition particularly in young children therefore it is important that interventions aimed at improving nutrition status of women and children are specific considering regional agricultural practices and seasonal differences.

This review examined undernutrition and its determinants based on the UNICEF conceptual framework on malnutrition. The underlying determinants including food security, environmental factors such as agricultural practices and seasons were also considered. Lastly, multicomponent interventions focusing on agricultural interventions and practices that are affecting immediate changes in nutrition status among infants and young children are emphasized.

2. Global Prevalence of Child Undernutrition

Globally, undernutrition affects a significant number of children below the age of five. An estimated 22.3% (148.1 million) of children are stunted and 6.5% are wasted in 2022 [164]. The rates of undernutrition vary worldwide with highest rates noted in the African and South-East Asia regions. UNICEF reported the prevalence of stunting in Africa at about 43% and wasting at about 27% These statistics can be broken down further by region with stunting rates in Southern Africa at 22.8%, followed by 30% in West Africa, East Africa at 30.6% and the highest recorded rates (37.4%) in Central Africa [164]. Within East-Africa, countries with the highest rates of stunting include Burundi 56.5%, Malawi 34%, Congo 40.3%, Ethiopia 34.4% and Tanzania 30.6% [164]. This shows that undernutrition continues to be a major public health problem in developing countries and that East Africa in particular has very high rates of chronic undernutrition.

Globally, micronutrient deficiencies (hidden hunger) are highly prevalent affecting 56% (372 million) among preschool-aged children, and 69% (1.2 billion) among

non-pregnant women of reproductive age. Regionally, three-quarters of preschool-aged children with micronutrient deficiencies live in south Asia (99 million) and sub-Saharan Africa (98 million) [148].

According to the World Health Organization (WHO), the most common nutrient deficiencies are vitamin A, folate, iron, iodine, zinc and Vitamin D [148] and the burdens of iodine deficiency and vitamin A deficiency remain high in Central Sub-Saharan Africa. Iron deficiency anaemia is the most common type of anaemia worldwide, which normally coexists with other deficiencies such as folic acid, vitamin B12 and vitamin C [61, 102]. According to WHO 40% of women of reproductive age of children below the age five years are anaemic [128, 149] and about 39% of children are vitamin A deficient [12]. The prevalence of anaemia, iodine and Vitamin A deficiency among under 5 children was reported to high in Southern Africa, East Africa, West and Central Africa [61] Vitamin A deficiency are persistently affecting children under 5 years of age in sub-Saharan Africa and this is due to insufficient dietary diversification, unsuccessful food fortification, and the restricted effect of vitamin A capsule supplementation on serum retinol [61].

3. Child Undernutrition in Tanzania

Tanzania has made progress in reducing child undernutrition from the past years according to the Tanzania Demographics Household Survey reports [114] (Table 1). There is a steady decline in child underweight and stunting from 1992 to 2022. Nevertheless, the prevalence of child underweight and stunting in 2022 was still high according to the criteria set by WHO [175]. About 35% of children below 5 years of age were stunted, 12% were underweight and 3% wasted ICF&MACRO, 2022. There are rural-urban and regional disparities in chronic undernutrition in Tanzania. In 2011, 33.4% of children in rural areas were stunted compared to 20.5% of children in urban areas. Despite a decrease in the proportions, there was a similar trend in 2015 with 45% of stunted children in rural Tanzania, compared to 32% in urban areas. In addition, to urban and rural differences there are differences between different regions in Tanzania (Table 2). According to WHO criteria to classify stunting prevalence, three regions had a very high severity of stunting (>40%), of 56.9, 49.8 and 50.4 recorded in Iringa, Rukwa and Njombe respectively [175]. In addition to the DHS reports, several regional studies have reported prevalence rates with highest rates of stunting (56%), underweight (46%) and wasting (24%) reported in Kilimanjaro region [106, 143] (Table 2).

Table 1. Nutrition status trends in Tanzania according to TDHS and TNNS.

Status	1992	1996	1999	2004/05	2010/11	2015/16	TNNS (2018)	2022
Stunting (%)	47	43	44	38	42	35	31.8	30
Underweight (%)	29	31	29	22	16	14	14.6	12
Wasting (%)	6	7	5	3	5	5	3.5	3

Source: Tanzania National bureau of statistics reports (NBS); Tanzania National Nutrition Survey (TNNS-2018)

Prevalence rates of micronutrient deficiencies are also high in Tanzania, particularly for children and pregnant women. Like stunting rates, the prevalence of anaemia mainly due to iron deficiencies has also declined from 2010-2016 (Figure 1). The prevalence of any anaemia ranged from ~85% in 2005 to 76% in 2016 with the highest rates among infants aged 9-11 months olds (81%) [114]. About 42% of children under age 2 years had moderate anaemia in 2016. This is a critical age during which undernutrition including essential

micronutrient deficiencies have irreversible lifelong impact on child health and development [12]. In addition to the age-related differences, there are also differences in prevalence between urban and rural children. There is slight variation in anaemia prevalence among rural and urban children, the prevalence of anaemia was slightly higher among rural children (73%) compared to urban children (67%) with rural children showing remarkable improvement in 2016 with the prevalence at 44.8% and urban 44.5% [114].

Table 2. Regional disparities in the prevalence of stunting, underweight and wasting.

Author	Region/District	Age group	Stunting	Underweight	Wasting
TDHS 2022	Geita	6-59mo	38.6	10.3	3.3
TDHS 2022	Njombe	6-59mo	50.4	12.2	2.1
TDHS 2022	Rukwa	6-59mo	49.8	19.4	8.3
TDHS 2022	Iringa	6-59mo	56.9	10.7	1.9
(Mrema et al., 2021)	Kilosa-Morogoro	6-59	41	11.5	2.5
Muhimbula et al., 2019	Shinyanga/Morogoro	0-24	40-42		
(Mgongo et al., 2017)	Kilimanjaro	0-24mo	42	46	24
(Munisi et al., 2016)	Rorya District		38	—	14
(Altare et al., 2016)	Southern highland	6-59mo	47	—	—
(Nordang et al., 2015)	Rukwa	6-59mo	64	34	—
(Kulwa et al., 2015)	Mpwapwa district	<1yrs	34	—	—
(Semali et al., 2015)	Kongwa district	< 5 yrs	49.7	—	—
(Shirima et al., 2015)	Tabora, Kilimanjaro and Iringa	<5yrs	56	—	—
(Chirande et al., 2015)	DHS data	0-5yrs	35.5	—	—
(Abubakar et al., 2012)	Kilimanjaro	1-35mo	44	19	5

Source: Author, 2023




Figure 1. Anaemia prevalence by age (months)-TDHS, 2005-2016.

4. UNICEF Framework for Undernutrition

The UNICEF conceptual framework remains the international preferred framework that illustrates the causes and consequences of malnutrition. It is a useful tool for understanding the determinants of undernutrition and serves as a guide in analysing the causes of undernutrition and planning interventions [158]. The framework explains the relationship between poverty as a root cause of undernutrition, food insecurity, among other immediate and underlying causes of maternal and child undernutrition (Figure 2). The framework has three levels of causality: immediate, underlying and basics, which should guide

interventions from a multi-sectoral viewpoint. The framework was developed further in the 2013 Lancet Nutrition Series (Figure.3) pointing out the determinants of nutritional status at different levels and outlining different sectors that are essential to respond effectively toward optimum fetal and child nutrition and development other than reducing undernutrition only [23].

4.1. The Immediate Causes of Undernutrition

The immediate causes occur at individual levels and are due to an imbalance between the nutrients absorbed by the body and nutrients required by the body due to inadequate food intake, poor nutrient utilization in the body or infections. Poor quality and quantity of food intake including inadequate intakes of protein, energy and essential micronutrients, lead to increased susceptibility to infections and undernutrition [23, 39]. In a vicious cycle, infections can cause poor appetite, and poor absorption of nutrients leading to reduced food intakes and hence undernutrition. The immediate causes affect children directly and can lead to short-term consequences, such as morbidity, mortality and disability, as well as long-term consequences, such as short adult stature, poor cognition and chronic diseases of a lifestyle (Figure 2). Infections such as malaria, diarrhoea, pneumonia and measles affect undernutrition by compromising immunity, leading to underweight and stunting [3, 30, 139]. In addition, poor maternal nutrition status before and during pregnancy has significant effect on child nutrition status therefore must be considered as part of the immediate impacts on child undernutrition [125].




Figure 2. UNICEF conceptual framework adapted from UNICEF 1990.




Figure 3. Framework for actions to achieve optimum fetal and child nutrition and development.

4.2. The Underlying Causes of Undernutrition

The underlying causes of undernutrition can be grouped under the three broad categories including; household food insecurity, inadequate care and feeding practices for children, and poor health and sanitation. Household food insecurity in terms of low availability and poor access was identified as an underlying cause of undernutrition in several studies [121, 137]. Inadequate care, poor health and sanitation was identified as causes of infectious diseases such as diarrhoeal diseases and respiratory infections increasing the risk of mortality [54, 82, 119]. Caring practices for young children and mothers during pregnancy and lactation with adequate nutrition and health, including proper medication, can lead to good nutrition. All these factors are underpinned by poverty due to lack of employment, remittances, and assets. Moreover, seasonal availability of food, agro-ecology and farming systems play major roles as underlying determinant of household food insecurity, mainly due to the impact on access and availability of food.

4.3. Basic Causes of Undernutrition

The basic causes of undernutrition are structures and processes around society that intensify poverty, particularly for vulnerable groups of society through limited and poor access to essential resources available. The complex interaction between socio-cultural, economic and political environment affect undernutrition in such a way that it impacts the environment in which individual lives (Figure 2). This can lead to poor financial, human and social capital mostly for women in the society. Poor socio-economic and political environments make it difficult for households to access land for farming, education, lack of employment, income and poor technology. However, high-level political commitment can make it easier for multisector actions to reduce undernutrition. Further, supportive policy environment across sectors are essential for improving nutrition by scaling-up of proven interventions as the key driver for changes. In order to attain food and nutrition security, national food and nutrition policies should increase access to services and credit, food safety, food market, on farm and off farm employment, irrigation and climate change mitigation [161]. All these rely on, government commitments to increase investment in these priority areas [116, 147].

For the sustainable impact on scaling up nutrition, there is a need for multiple sectors in agriculture, public health, water, sanitation and hygiene and social protection working together to combat undernutrition; [55]. The framework (Figure 3) shows this multisectorality and the multi-level nature of responses. It explains the nutrition-specific interventions (delivered by the health sector), and nutrition-sensitive interventions (delivered by other sectors), both depending on enabling policy environments that are needed to provide support for nutrition programmes [23]. Nutrition-sensitive programs in agriculture, social safety nets, early child development and education have potential, to augment the

coverage and effectiveness of nutrition-specific actions although its potentials have not been adequately unleashed [131]. Additionally, nutrition-sensitive interventions can create a stimulating and health environment for optimum fetal and child growth and developments to their full potential (Figure.3).

5. WHO Infant and Young Child Feeding (IYCF) Practices

UNICEF and other child nutrition and health experts agree that optimal IYCF has the single greatest potential impact on child survival, compared to all proven preventive health and nutrition interventions [26, 159, 177]. Therefore, reduction of child undernutrition, morbidity and mortality can be reached when nutrition in early infancy and IYCF specifically are highly prioritized in national policies and programme strategies. The benefits of breastfeeding and optimal complementary feeding for child survival, growth, and development are well known and are estimated to prevent 823,000 child annual deaths in low-income and middle-income countries [166]. Studies show that exclusive breastfeeding is the best form of nutrition for children up to 6 months, the composition is dynamic and personalised for each child; it lowers infectious morbidity and increases intelligence and cognition, persisting to adulthood [166]. Appropriate complementary feeding practices are important to maintain a child's growth potential when requirements for energy, protein and other nutrients (particularly iron and zinc) cannot be met by breast milk alone from the age of 6 months [115]. If appropriate complementary feeding practices could be scaled up to nearly universal levels, ~100,000 deaths in children under five could be prevented each year [27]. Additionally, the benefit of optimal complementary feeding practices as a preventive measure of stunting among 6-24 months has been as important [150]. Therefore, complementary foods should be of high nutrients density due to increased nutrients requirements for metabolic processes, rapid developmental processes, and limited gastric capacity for infants from 6-24 months failure to deliver on this guideline which leads to growth faltering [39, 58]. The next section describes the WHO IYCF indicators and recommendations and discuss the relationships between feeding practices and nutrition outcomes.

5.1. IYCF Indicators and Adherence to Recommendations

WHO/UNICEF developed simple and valid age appropriate indicators to assess breastfeeding and complementary feeding practices among children less than 2 years old which are amenable to population-level measurement [176]. There are 8 core indicators and 7 optional indicators used as means of monitoring adherence to IYCF recommendations worldwide (Table 3).

Table 3. WHO-recommended core indicators for measuring IYCF practices (WHO, 2008).

Core indicators	Description of indicator
1. Early initiation of breastfeeding*	Proportion of children born in the last 24 months, breastfed within 1 hour of delivery.
2. Exclusive breastfeeding under 6 months	Proportion of infants 0–5 months of age who received only breast milk during the previous day.
3. Continued breastfeeding at 1 year	Proportion of children 12–15 months of age who received breast milk during the previous day.
4. Introduction of solid, semi-solid or soft foods	Infants 6–8 months of age who received solid, semi-solid or soft foods during the previous day.
5. Minimum dietary diversity	Proportion of children 6–23 months of age who received foods from 4 food groups during the previous day.
6. Minimum meal frequency	Proportion of breastfed of children 6–23 months of age who received solid, semi-solid or soft foods the minimum number of times or more during the previous day.
7. Minimum acceptable diet	Proportion of breastfed of children 6–23 months of age who received at least the minimum dietary diversity and the minimum meal frequency on the previous day (apart from breast milk).
8. Consumption of iron-rich or iron-fortified food	Proportion of children 6–23 months of age who received an iron-rich food or an iron-fortified food on the previous day.
Optional indicators	Description of indicator
Ever breastfed	Proportion of children born in the last 24 months who were ever breastfed.
Continued breastfeeding at 2 years	Proportion of children 20–23 months of age who are fed breast milk.
Age-appropriate breastfeeding	Proportion of children 0–23 months of age who are fed according to guidelines.
Predominant breastfeeding under 6 months	Infants 0–5 months of age who received breast milk as the predominant source of nourishment during the previous day
Duration of breastfeeding	The median age in months when 50% of children 0–35 months did not receive breast milk during the previous day.
Bottle feeding	Proportion of children 0–23 months of age who are fed with a bottle during previous day.
Milk feeding frequency for non-breastfed children:	Proportion of non-breastfed children 6–23 months of age who receive at least 2 milk feedings during the previous day

Source: Indicators for assessing infant and young child feeding practices, conclusions of a consensus meeting held 6–8 November 2007 in Washington DC, USA (Working Group on Infant and Young Child Feeding Indicators 2007. [176])

*Provision of mother's breast milk to infants within one hour of birth is referred as "early initiation of breastfeeding"

Globally, 47% of infant are put to the breast within 1 hour of delivery and only 43% are exclusively breastfed 0-5 months [178]. Breastfeeding is commonly practiced in developing countries and nearly all infants are put to the breast, however, sub-optimal adherences to recommended practices are common. According to UNICEF statistics, the prevalence of early initiation in Eastern and Southern Africa was 65% and exclusive breastfeeding for 0-5months was 55% In [178]. Tanzania, almost 99% of infants are breastfed, regardless of socio- economic categories; however, early initiation and exclusive breastfeeding are persistently suboptimal (Table 4). The proportions of initiation of

breastfeeding have not changed considerably since 1996 and it shows decline in 2010. There is improvement with exclusively breastfeeding 0-5 yearly but the target has not been reached, more than 40% of infants are not exclusively breastfed. This indicates that a substantial proportion of infants 0-5 months are given other fluids and solid foods before the recommended age of 6 months, which hinders the benefit of breastmilk to the majority [166]. Additionally, there are regional disparities and rural urban differences with regards to exclusive breastfeeding practices where high rates and very low rates were reported, which need to be considered when planning interventions [114].

Table 4. Trends in initiation and exclusive breastfeeding in Tanzania.

Months	Initiation	0-1	2-3	4-5	6-7	0-5
Year	%	%	%	%	%	%
1996	59.0	55.2	27.4	8.0	4.1	28.9
1999	NA	57.8	25.4	15.5	1.9	NA
2005	59.3	70.0	42.4	13.5	1.7	41.3
2010	46.1	80.5	51.1	22.9	2.3	49.8
2015-16	51.2	84.0	58.8	26.6	3.0	59.0

Source: Tanzania National Bureau of Statistics (NBS), Data from Demographic and health survey (TDHS)

Globally, only 29% of infants are fed from at least 4 food groups and 16% have a minimum adequate diet [162]. About 21% of infants aged 6-23 months had met minimum dietary diversity and only 10% met minimum adequate diet in Eastern and Southern Africa [162]. In Tanzania, 97% infants 6-8 months received solid foods which ties in with early introduction of solid foods, which is a common practise over the years [109, 114]. In addition, only 8% received a

minimum acceptable diet and less than 40% had minimum meal frequency [114]. Low adherences to complementary feeding recommendations were observed among rural children (7%) compared to urban children (12%), with 39% of urban children and 19% of rural children meeting minimum dietary diversity. [114]. There are limited published studies in Tanzania that reported on complementary feeding practices per WHO IYCF guidelines

[176]. One local study revealed poor compliance to complementary feeding recommendations [99] (Table 5).

The characteristics of the complementary diet in developing countries are widely reported, however, there are limited published data on nutrient contents of complementary foods in Tanzania. The diet is characterized by low intakes of animal and dairy products and higher intakes of cereal based porridge and legumes [1, 6, 39]. Foods consumed have low nutrient density, are unlikely to meet the requirements for

energy, protein and fat and key nutrients for growth particularly iron, zinc and calcium, vitamin A, vitamin B12 and niacin. Ideally, complementary foods should supply up to 95% of iron, 55% riboflavin, 60% calcium, 70% thiamine, 85% niacin and 85% vitamin B6 (Allen, 2012). Given the low intakes of costly animal source foods in the diet of infants, it is challenging to meet nutrient needs in the critical period for growth without including animal source food [39, 6, 1].

Table 5. Adherence to IYCF recommendations.

Year	Age (months)	DDS (4+ food groups)	Minimum meal frequency	Minimum acceptable diet
2010*	6-8	41.6	64.6	27.7
	9-11	57.7	24.9	18.0
	12-17	69.5	32.5	23.1
	18-23	68.0	38.5	29.1
Urban		68.1	42.4	27.1
Rural		59.5	37.6	23.6
2015-16	6-8	11.3	66.0	9.3
	9-11	24.0	35.8	8.4
	12-17	29.9	38.0	12.1
	18-23	25.6	34.7	7.0
Urban		36.4	37.3	13.2
Rural		19.6	44.9	8.6
Kulwa et al., 2015 (Rural)	6-11	4.6		
Vital et al., 2016 (Urban)	6-23	49.3	70.3	38.4
Victor et al., 2014	6-23	38.2	38.6	15.9
(Masuke et al., 2021)	6-24		40.4	

For breastfed children, minimum meal frequency is receiving solid or semi-solid food at least twice a day for infants age 6-8 months and at least three times a day for children age 9-23 months. * TDHS 2010 used 3+ food groups for dietary diversity indicator: DDS=Dietary diversity score

Source: Author(s); Tanzania National Bureau of Statistics (NBS)

5.2. Factors Associated with Meeting IYCF Recommendations

The global strategy on infant and young child nutrition (IYCN) highlighted that inadequate knowledge regarding proper feeding practices and cultural beliefs are major constraints to successful feeding practices other than food availability [163]. Studies have reported that social and cultural factors influence a mothers adherence with IYCF guidelines, including; 1) mothers perceive their infant to be “thirsty” and believe that breast milk contains insufficient water; 2) mothers believe that breastfeeding should be stopped immediately when one becomes pregnant again as the milk of a pregnant mother can harm the infant; and 3) mothers fear that breastfeeding too long will lead to the infant to becoming addicted to breast milk, which will cause them rejecting solid food at the time of introduction [34]. Other studies have reported inadequate breastfeeding skills, peer pressure and influences of other family members as major constraints to successful breastfeeding [80]. Furthermore, mothers’ beliefs that they have insufficient milk to feed infants exclusively up to 6 months, HIV status hindering mothers to choose breastfeeding as an option and maternal employment forcing mothers to leave the child early for work/business were also reported [34, 79, 90]. On the other hand, mothers who delivered at the hospital were likely to practice early initiation compared to home delivery [42].

Lower maternal education, young maternal age and lack of professional assistance at birth where also determinants to poor feeding practices [63, 76, 165] Household food insecurity was identified as a determinant of poor complementary feeding practices and also mothers who experienced household hunger had early cessation of exclusive breastfeeding and believed that they need adequate foods to breastfeed up to 6 months [37, 172, 181]. This means that education interventions for proper feeding practices are not reaching most mothers/caregivers. Comparing to urban, rural mothers have poor breastfeeding and complementary feeding practices in Tanzania [114]. The rural-urban differences are partly explained by insufficient resource allocation where families in rural areas have poor access to health services, clean water, nutrient-dense foods and nutrition information.

5.3. IYCF Practices and Child Nutrition Outcomes

Age appropriate IYCF practices are some of the major determinants of good nutrition status and growth among infants and young children [20]. For example, early initiation of breastfeeding was reported to lower the risk of underweight and stunting [64, 93, 92]. In addition, exclusively breastfeeding was associated with reduced risk of underweight and wasting [85,93]. However, some studies found no relationship with early initiation of breastfeeding and exclusive breastfeeding[40,104]. The possible

explanation of the relationship between proper breastfeeding practices and weight gain could be related to more frequent breastfeeding at a young age enabling infants to get enough breast milk and adequate nutrients for growth. On the contrary, continued breast feeding at 12-15 months was linked to higher risk of underweight [93, 85]. This could be related to inadequate complementary feeding practices with poor quality and fewer meal frequencies, the relationship that has been reported earlier [49, 95].

Timely consumption of solid foods at 6-8 months and proper meal frequency was reported to lower the risk of underweight [92] and was associated with higher WAZ and WHZ [135, 157, 182] and higher LAZ [40]. Proper meal frequency is associated with adequate energy and the type of solid foods given to infants could be related to infant weight gain or weight loss. However in many developing countries, usually infants start with thin maize porridge, low in energy and with low intakes of vitamin A-rich and animal source foods which could be related to poor weight gain. This is supported with findings in Sri Lanka where the late introduction of fat and oils into the diet of infants after 8 months of age was associated with underweight and wasting indicating the importance of timely introduction of a high quality diet for proper weight gain [156]. Finally, the relationship between diet diversity and nutrition status has been reported frequently; for example, diet diversity score is associated with a reduced risk of stunting [9, 36, 43, 56, 88, 89, 124, 130]. Both dietary diversity and the consumption of iron rich foods were associated with a reduced risk of both underweight and stunting [64, 85, 92]. Furthermore, dietary diversity correlated with adequate energy intakes [9,14] and improved haemoglobin concentrations particularly if animal source foods were included [81, 134]. This shows that dietary diversity is an indicator of diet quality [132].

6. Maternal Nutrition Status

Women's health and wellbeing depend on the complex interaction of a number of different factors including, social and cultural environment, access to health services, and life-stage [107]. The foetus depends on the mother throughout pregnancy for their nutrient supply, therefore maternal nutrition status is a good indicator of her own health and reproductive ability and is a good predictor of new born health and nutrition status [154]. Research suggests that poor nutrition before and during pregnancy accounts for 50% of infant growth failure, which has long-term implications for the child into adulthood [160, 25]. Two major concerns in Tanzania are thinness, and anaemia. Thinness (BMI <17 kg/m²) affects 10% of women of reproductive age and about 57% of pregnant women are anaemic [114]. Acute nutrition deficiencies during pregnancy increase the risk of abnormal delivery and lowers immunity of both mother and child [91]. Improving nutrition status and health of women and children is a priority area for action [20].

Maternal Nutrition Status and Child Nutrition Outcomes

Poor maternal nutrition status impacts foetal growth

leading to increased risk of small for gestational age births (SGA), LBW and preterm birth [20-22]. Infants born SGA or with LBW are more likely to be stunted, have a higher risk of poor-quality life and poor cognitive outcomes (Prendergast and Humphrey, 2014). Moreover, LBW is the biggest risk factors for mortality and morbidity in the first few months of life, particularly in developing countries [127]. Poor weight gain during pregnancy due to inadequate dietary intakes and maternal workload are known risk factor for LBW (Beilly & Kurland, 1945; [18, 22, 35, 87, 100]. Conversely, higher energy and protein intakes, and maternal haemoglobin status are associated with normal birthweight [105, 153, 155].

Other research has looked at the relationship between maternal anthropometrics post-partum and infant nutrition status. Acknowledging that the relationship between maternal height and child height is partly due to genetics [154]; in resource, poor settings where food insecurity and nutrition are major concerns there are several other factors that influence maternal height and weight and the relationship with infant anthropometrics is clear. Studies report relationships between maternal BMI and height, and higher WHZ scores and lower risk of stunting and wasting [104]. Another study reported that mothers' diet during pregnancy was related to her weight gain during pregnancy and subsequently the child's weight and height [53]. Altogether, this gives promising results in fighting undernutrition through investing in mothers and addressing the devastating impact of undernutrition during the first 1000 days. More emphasis on nutrition intervention from adolescence through early pregnancy and lactation is therefore needed. It is important to address maternal feeding and health during pregnancy and lactation and to identify mothers with poor weight gain during pregnancy and provide support where possible.

7. Food and Nutrition Security

Food and nutrition security (FNS) combine two concepts, food security and nutrition security. Adopted from the 1996 World Food Summit "food security exists when all people at all times have physical, social and economic access to sufficient, safe and nutritious food to meet their dietary needs and food preferences for an active and healthy life" [44, 45]. Unlike food security, nutrition security (built from the UNICEF conceptual framework) is defined as "having a nutritionally adequate diet that is biologically utilized to maintain growth, resisting or recovering from disease, pregnancy, lactation and physical work" [51]. Food and nutrition security is in turn defined as a condition under which adequate food is available and accessible and satisfactorily utilized by all individuals at all times to live a healthy and happy life [173].

Food Security and Child Nutrition Outcomes

Food security exists when all people always have physical and economic access to sufficient, safe and nutritious food to meet their dietary needs and food preferences for an active and healthy life [45, 46]. Food insecurity at global, national

and household levels has devastating effects on health and human development which is aggravated by climate changes [47]. Despite the progress made in increasing world food production, 3.1 billion people are unable to afford a healthy diet globally [48]. Persistent undernutrition (particularly hidden hunger) in Sub-Saharan Africa are due to food insecurity and poor diet quality and it affect 220 million people [78]. Moreover, changes in global environments, weather pattern and climate variability have impacted food production and food access for subsistence farmers, increasing the risk of food insecurity, infectious disease and undernutrition particularly for young children [48, 126].

Food insecurity, hidden hunger and undernutrition are still challenging in Tanzania, underpinned by seasonal and regional food shortages. The United Republic of Tanzania ranked 160 out of 191 (Human Development Index (2022) and 95th out of 121 on the 2022 Global Hunger Index with scores 23.6 which indicate serious population undernourishment [170]. The majority of food insecure households live in rural areas where agriculture is major source of livelihood. They rely on subsistence farming which is vulnerable to climatic, economic and seasonal shocks and they are battling hunger [74, 174].

Informing policy making and planning for interventions for child undernutrition is suitable but requires better understanding of several factors that influences children's health and nutrition. The relationship between household's food insecurity and certain nutrition status parameters among children below five years of age has widely been established and yielded mixed results. It is known that children from food insecure households can have lower food consumption, including fewer numbers of food groups, poor nutrient intakes and nutritional status [110]. However, other research shows that within food insecure households, children are better nourished because available foods will be given to young children as one of the coping strategies, while others argue that in food insecure households' young children receive more benefit from breast milk because of reduced food availability [57].

For example, in Nepal, children from severely food insecure households were highly stunted and underweight compared to children from food secure households, however, the rate of undernutrition among children from food-secure households was also high [145]. This indicates other determinants of poor nutrition than food insecurity, which could be related to poor feeding and care practices, frequent infection, poor sanitation, poor nutritional knowledge leading to nutrition deterioration. Similarly, within food insecure households children were more underweight, stunted but not wasted compared to food secure counterparts [4, 17, 145]. Wasting is an indicator of hunger or food/nutrition insecurity and low food intakes, while stunting is the results of chronic food shortage, poverty, infections and overall low social economic status. The impact of food insecurity on height and weight in the reported studies could indicate multiple problems in food insecure households. Some researchers reported underweight among children from food insecure

households [110, 134-135] whereas other reported lower risk of stunting among infants from food secure households [113, 123, 133]. Both stunting and underweight were reported in food insecure households [136]. On the contrary, other studies found no relationship between households' food insecurity and child anthropometry, only households wealth index was associated with undernutrition among children aged 6 to 23 months [121]. Some studies reported dietary diversity as a good predictor of stunting compared to household food insecurity particularly in communities where dietary diversity is high. It is therefore likely that the indicator is a stronger predictor of stunting than household food security.

8. Agriculture and Livelihoods in Africa

Worldwide, the livelihood of 2.5 billion people particularly small-scale farmers, herders, fishers depends on agriculture [47]. Africa mainly Sub-Saharan Africa is still facing great challenges of food insecurity with relatively low levels of agricultural productivity, low incomes among rural farmers and, high rates of undernutrition (AGRA, 2016). Agriculture is linked to nutrition particularly among subsistence farmers in developing countries as a source of income for food and non-food expenditures which is possible through employment in agricultural sector and selling agricultural produce, which can be translated into expenditure on nutrition-enhancing goods and services [55]. Therefore, agriculture is an important sector, critical to the economies of all African countries, where about 70% of population dependent on agriculture for livelihoods with possibilities of eradicating poverty and hunger (NEPAD, 2013). At the African Union Summit in Malabo, Equatorial Guinea in June 2014, Heads of State and Government adopted a remarkable set of concrete agriculture goals to be achieved by 2025. The Malabo Declaration on Accelerated Agricultural Growth and Transformation for Shared Prosperity and Improved Livelihoods is a set of new targeted approach to achieve the agricultural vision for the continent which is shared prosperity and improved livelihoods. The member state was commitment to Zero hunger – Ending Hunger by 2025 by at least double agricultural productivity by improving irrigation and mechanization and reducing post-harvest losses. The Malabo declaration set ambitious targets for African children there aiming to improve children's nutrition status by reducing stunting to 10 percent and underweight to just 5 percent by 2025 with agriculture as a key strategy. The Malabo Summit reconfirmed that "agriculture should remain high on the development agenda of the African continent and is a critical policy initiative for African economic growth and poverty reduction" (Malabo Declaration, 2014).

8.1. Agriculture and Season

Agricultural production depends on weather patterns throughout the year. In subsistence farming systems where food consumption is dependent on production, seasons and

rainfall, changes in weather patterns have an impact on production and therefore food consumption (Kaminski et al., 2016). The pre-harvest season is characterized by rainfall, intensive agricultural labour and low food stocks among all the farming households, affecting food security and livelihoods (Ali et al., 2016). Recently, climate change has had an impact on seasonal stress among rural farming households affecting quantity and types of food grown, nutrition status and lack of water [5]. Over the past number of years there have been an increased focus on the relationship between agriculture production and nutrition [59,103]. The attention on agriculture was emphasised in the recent Zero Hunger Challenge [152] and Sustainable Development Goals (SDG), which aims to “end hunger, achieve food and nutrition security and promote sustainable agriculture” [86, 97].

8.2. Agriculture and Child Nutrition Status

Agriculture is the primary source of livelihoods for food and income and is a strong driver of food and nutrition security [142]. It is therefore imperative to consider agriculture production and child nutrition through its influence on dietary intakes and feeding practices as immediate causes of undernutrition, household food insecurity and income as underlying determinants of nutrition status [55]. The pathway in which agriculture is linked to food security and nutrition outcomes is summarized below [41, 55]

1. Agriculture is a source of food (nutrients rich) for households' consumption: this is the most direct pathway whereby household increase food availability and accessibility from own production.
2. Agriculture provides a source of income for food and non-food expenditures: income is generated through employment in agricultural sector and/or selling agricultural produce, which can be translated into expenditure on nutrition-enhancing goods and services (including health, education, and social services).
3. Agriculture affect food price: effects of agriculture policy and food prices on food consumption, which in turn affect purchasing power of net buyers particularly rural subsistence farmers if the price of is high.
4. Effects of women's employment in agriculture on intra-household decision-making and resource allocation: agricultural labour conditions can influence the empowerment of women and thus their control over nutrition-relevant resources and decision making, particularly regarding food and healthcare.
5. Effects of women's employment in agriculture over time allocation: time allocation can challenge women to balance time spent in income generating activities, household management and maintenance, caregiving, and their own leisure
6. Effects on agricultural labour on women's own nutritional and health status: relates to the energy intensive nature of agricultural labour and effects on maternal nutrition and health status including high

energy expenditure and exposure to agriculture-associated diseases through zoonotic diseases and waste water irrigation

Although the agriculture-nutrition pathways overlap, the first pathway (agriculture as a source of food (nutrients rich) for households' consumption is important for rural agricultural dependent households due to consumption of own production with its influence on child nutrition [29, 75]. Although there is limited evidence for a link between agriculture and nutrition, particularly with children [98, 131]. Carleto and colleagues reviewed some evidences on the link between livestock farming, dietary diversity and nutrition status of children [29, 117]. The link between agriculture and nutrition outcomes depends on the type of farming practices. Nutrition sensitive farming practices are likely to influence positive nutrition outcomes; for example, β -carotene-rich orange sweet potato was shown to improve vitamin A status of pre-school children in Mozambique and Uganda [70,122]. Integrated homestead food production coupled with a behaviour change communication programme, targeting women and children documented positive effects on child nutrition outcomes [120]. Other research found a positive relationship between agriculture production diversity and dietary diversity among children aged 6–23 months and higher LAZ among older children aged 24–59 months in Zambia [31,84]. In addition, livestock farming impacted nutrition outcomes of young children; for example, cow ownership and child milk consumption was associated with higher LAZ and reduced odds of stunting among children aged 6–24 months in Ethiopia (Hetherington et al., 2017; Hoddinott et al., 2015). On the other hand, agricultural interventions to increase vegetables and fruits showed less impact on nutrition status, however livestock ownership and milk consumption were associated with improved WAZ and HAZ among children below the age of 5 years in Kenya [171]. The ownership of small ruminants was linked to lower wasting and underweight among 24–59 months old in Uganda [11]. Additionally, crop output value had small effects on anthropometric outcomes of pre-schoolers and children 5–9 years old while livestock ownership had positive outcome on the nutrition of children in Tanzania [146]. Few studies have confirmed the strength of agriculture production and food consumption to effect changes among infants and young children 6-24moths in rural farming households.

8.3. Season and Child Nutrition Status

Seasonal fluctuations in food availability is an important factor influencing child growth, particularly in rural farming households where season impacts food availability (Wijesinha-Bettoni et al., 2013). The effect of season on food availability and child anthropometrics has been observed in literature, with studies reporting better nutrition during the post-harvest season [50, 62, 67]. The typical drop in food stock prior to the harvest season results in a decrease in food consumption. At the same time, the time dedicated to farming increases, which has an impact on energy

expenditure. The combination of the drop-in food availability and the increased workload means a mismatch in energy balance and a higher risk of undernutrition [32, 68]. In line with decreased food consumption, incomes and food prices change with the agricultural seasons which puts additional pressure on vulnerable groups [68]. In addition, seasonal weather fluctuations affect food availability and environmental sanitation and health thereby affecting diet, infant feeding and morbidity [15, 118, 179]. Pre-harvest seasons, lower WAZ and WLZ was reported among children in farming households [72]. A decrease in wasting and underweight was reported during post-harvest in Ethiopia indicating increased food intakes during this time [129]. On the contrary, seasonal morbidity was a significant cause of seasonal fluctuation in nutrition status among children under the age of five years in Malawi compared to food availability [32]. Some authors have linked variation in child nutrition status across agricultural seasons as lack of care from mothers due to high agricultural labour during pre-harvest and post-harvest and not necessarily with food availability [118,151]. From the documented studies, the link between seasonal food availability, dietary intake and infant feeding practices among infants and young children is lacking, and it is not clear if variation in feeding practices indicators, nutrients intakes and nutrition status follow seasonal patterns of food availability, creating the necessity of examining children's diets at different times of the year.

9. Targeting Improvements in Child Nutrition Status

The analysis of early growth patterns in children from resource-poor countries shows a rapid decline in the Length for age Z-score during the first 2 years of life with no recovery until 5 years. This observation brought attention to the first years of life as a window of opportunity for targeting interventions against stunting [144] (Victora et al., 2010). Although some studies have presented other windows of opportunity for intervention to curb stunting outside the 1000 days windows (Prentice et al., 2013), the adverse effect of undernutrition during this period was established [21-22]. Women particularly in low-income countries enter pregnancy when they are young and in a state of poor nutrition. The additional demands of pregnancy increase the risk for undernutrition for these women and their infants [96]. This chronic, inter-generational state of undernutrition is hampering the potential of future generations in developing countries. Dietary diversification and consumption of nutrient-rich foods among women, coupled with proper feeding practices for infants and young children is the primary long-term goal for improving nutrition during the target periods [39]. Policy-makers need to focus on multicomponent interventions that target the immediate and underlying causes of undernutrition in order to affect long-term change in nutrition status outcomes in developing countries [77, 168].

10. Multi-Component Interventions to Improve Child Nutrition Status

There are various determinants of undernutrition reviewed prior to this section (figure 2), however, lack of connection between different sectors including agriculture, public health, education, water and sanitation, social protection, individual and household decision making lead to poor improvement in child nutrition [60]. Therefore, multicomponent interventions are necessary to improve child nutrition status in the long term. This has been analysed in figure 3 addressing the multisector responses to fight undernutrition [26]. Acknowledging the importance of adequate nutrition during the formative years has caused tremendous increases in political commitment for the reduction of under-nutrition at the global and national level, focusing on the first 1000 days of life. This commitment includes the global effort on improving nutrition through agriculture particularly targeting the first 1000 days [131, 138]. This was emphasised through the Sustainable Development Goals which aims of “end hunger, achieve food and nutrition security and promote sustainable agriculture” [86]. While agriculture is a key underlying sector, its effectiveness depends on collaboration with other sectors like education, water and sanitation, social protection, early child development and nutrition. This has led to the reorientation of many programs to focus on nutrition sensitive interventions to address the underlying determinants of foetal and child nutrition. Nutrition-sensitive programs in agriculture, social safety nets, early child development and education have potential, to augment the coverage and effectiveness of nutrition-specific actions although its potentials have not been adequately unleashed [131]. Additionally, nutrition-sensitive interventions can create a stimulating environment for optimum foetal and child growth and developments to their full potential (Figure 3). Agricultural intervention particularly, has a potential to improve livelihoods, food security and diet quality and can reach vulnerable population in large scale [181].

For a long time, in Tanzania, nutrition specific interventions (aimed at the immediate causes) have been in place addressing both maternal and child nutrition (Table 6). These include health interventions that impact nutrition outcomes such as iron and folic acid supplementation for pregnant mothers, vitamin A supplementation deworming and promotion of exclusive breastfeeding. However, the coverage of nutrition-specific interventions is low, indicating the needs for multicomponent interventions. For example, interventions based on nutrition sensitive agriculture, health, social protection, education, and water and sanitation including poverty reduction, can be implemented at large scale and can effectively reach poor populations and vulnerable groups [41, 132]. Additionally, policies and programmes that improve agriculture, health, education, social protection, water, sanitation and hygiene reduce stunting by 80%, compared to a 20% reduction with direct nutrition interventions [73]. Accordingly, it is important to focus on agriculture as a key sector in producing healthy and

nutritious diets that address both immediate and underlying determinants of undernutrition through coverage of large rural poor populations who depend on agriculture for food and livelihoods.

Despite strong evidence to support links between

agriculture, growth and a reduction in under-nutrition [29, 55] there is limited evidence to show the impact of nutrition sensitive agricultural interventions on child nutrition outcomes in Tanzania. This is calling out more research on nutrition farming practices and child nutrition outcome.

Table 6. Nutrition specific interventions in Tanzania targeting young children.

Type of interventions	Demographic & Health surveys	
	2010 (%) *	2015-16 (%)
Exclusive breastfeeding (0-5months)	49.8	59.2
Complementary feeding with at least 4 groups per day (6-23 months)	56.4	23.9
Zinc treatment for diarrhoea	4.7	17.5
Pregnant women attending 4 or more ANC visits	42.7	50.6
De-worming (12-59 months)	49.6	37.6
Vitamin A supplementation (6-59 months)	60.8	41.2
Households with adequately iodised salt (15+ ppm)	55.2	60.6
Postpartum women with Vitamin A supplementation	25.8	–
Insecticide-treated nets	73.9	54.4
Malaria prophylaxis for pregnant women	26.3	34.6
Folic and iron supplementation	40.0	21.0
Deworming for mothers	–	63.1

TDHS 2010 used 3 food groups for DDS, ANC: Antenatal care ANC=Antenatal care
Sources: Tanzania National Bureau of Statistics: TDHS 2010 and TDHS 2015-16

In conclusion, undernutrition remains a major global problem and an underlying cause child deaths annually. Proper nutrition is critical for child growth and survival particularly in infancy and early childhood when failure to grow can have lifetime consequences. Therefore, the “first 1000 days” is the ideal target for interventions aimed at reducing all forms of undernutrition in children. Efforts to reduce undernutrition in Tanzania have been largely focused on healthcare, including measures such as vaccination programmes, nutrition rehabilitation and micronutrient supplementation, which reflect Western medical practices and the medical model. Agricultural, environmental and social factors, which underpin the root cause of undernutrition have not been given much attention. Focusing on agriculture to improve nutrition, particularly targeting the “first 1000 days”, moving away from the traditional single nutrient approach to a food-based, comprehensive approach is the global emphasis globally [8, 55, 180]. Rural vulnerable households in Tanzania are an ideal target for this type of intervention given the high rates of undernutrition, poor breastfeeding and complementary feeding practices and high dependency on agriculture for livelihoods. Although we recognize the relationship between agriculture and nutrition, we need to identify targets before we can design appropriate interventions. In addition, Tanzania is an agriculturally diverse country with substantial regional variation in the prevalence of undernutrition. It is important that interventions aimed at improving nutrition status of women and children are specific and consider regional agricultural practices and the impact of season. While previous studies reported the determinants of child undernutrition, very few studies have taken a comprehensive approach including factors from the environment right through to infant feeding practice. This information is important to provide the evidence-base to inform policy-making and practice. One of

the key questions that remain. Does agriculture have the strength to affect immediate change in nutrition status during the first 1000 days? In an attempt to answer this question, the need to study the immediate, underlying and environmental factors that determine undernutrition among infants and young children and identified nutrition sensitive agricultural practices that can potentially form the basis of future intervention targets is warranted.

Conflicts of Interest

The author declares no conflicts of interest.

References

- [1] Abeshu, M. A., Lelisa, A., & Geleta, B. (2016). Complementary Feeding: Review of Recommendations, Feeding Practices, and Adequacy of Homemade Complementary Food Preparations in Developing Countries—Lessons from Ethiopia. *Frontiers in Nutrition*, 3.
- [2] Abubakar, A., Uriyo, J., Msuya, S. E., Swai, M., & Stray-Pedersen, B. (2012). Prevalence and risk factors for poor nutritional status among children in the Kilimanjaro region of Tanzania. *International journal of environmental research and public health*, 9(10), 3506-3518.
- [3] Alexandre, M. A. A., Benzecry, S. G., Siqueira, A. M., Vitor-Silva, S., Melo, G. C., Monteiro, W. M.,... Alecrim, M. d. G. C. (2015). The association between nutritional status and malaria in children from a rural community in the Amazonian Region: a longitudinal study. *PLoS Neglected Tropical Disease*, 9(4), e0003743.
- [4] Ali Naser, I., Jalil, R., Muda, W., Manan, W., Nik, W., Suriati, W.,... Abdullah, M. R. (2014). Association between household food insecurity and nutritional outcomes among children in Northeastern of Peninsular Malaysia. *Nutrition research and practice*, 8(3), 304-311.

- [5] Ali, S., Zareba, F. E., Maruod, M. E., Elkhidir, E. E., & Mahmoud, T. E. (2016, 4 – 12 March). Climate Variability and Change: Implications for Household Food Security in North Kordofan State. Proceedings of the Summer-School Workshop, University of Western Kordofan; Sudan.
- [6] Allen, L. H. (2012). Global dietary patterns and diets in childhood: implications for health outcomes. *Annals of Nutrition and Metabolism*, 61(Suppl. 1), 29-37.
- [7] Altare, C., Delbiso, T. D., Mutwiri, G. M., Kopplow, R., & Guha-Sapir, D. (2016). Factors associated with stunting among pre-school children in southern highlands of Tanzania. *Journal of Tropical Pediatrics*, 62(5), 390-408.
- [8] Ambikapathi, R., Schneider, K. R., Davis, B., Herrero, M., Winters, P., & Fanzo, J. C. (2022). Global food systems transitions have enabled affordable diets but had less favourable outcomes for nutrition, environmental health, inclusion and equity. *Nature Food*, 3(9), 764-779.
- [9] Arimond, M., & Ruel, M. T. (2004). Dietary diversity is associated with child nutritional status: evidence from 11 demographic and health surveys. *The Journal of nutrition*, 134(10), 2579-2585.
- [10] Arimond, M., Wiesmann, D., Becquey, E., Carriquiry, A., Daniels, M., Deitchler, M.,... Kennedy, G. (2011). Dietary diversity as a measure of the micronutrient adequacy of women's diets in resource-poor areas: summary of results from five sites. *Washington, DC: FANTA-2 Bridge, FHI*, 360, 2011.
- [11] Azzarri, C., Zezza, A., Haile, B., & Cross, E. (2015). Does livestock ownership affect animal source foods consumption and child nutritional status? Evidence from rural Uganda. *The Journal of Development Studies*, 51(8), 1034-1059.
- [12] Bailey, R. L., West, K. P., & Black, R. E. (2015). The Epidemiology of Global Micronutrient Deficiencies. *Annals of Nutrition and Metabolism*, 66, 22-33. <https://doi.org/10.1159/000371618>
- [13] Bain, L. E., Awah, P. K., Geraldine, N., Kindong, N. P., Siga, Y., Bernard, N., & Tanjeko, A. T. (2013). Malnutrition in Sub-Saharan Africa: burden, causes and prospects. *Pan African Medical Journal*, 15(1).
- [14] Bandoh, D. A., & Kenu, E. (2017). Dietary diversity and nutritional adequacy of under-fives in a fishing community in the central region of Ghana. *BMC Nutrition*, 3(1), 2.
- [15] Barrett, B., Charles, J. W., & Temte, J. L. (2015). Climate change, human health, and epidemiological transition. *Preventive medicine*, 70, 69-75.
- [16] Beilly, J. S., & Kurland, I. I. (1945). Relationship of maternal weight gain and weight of newborn infant. *American Journal of Obstetrics and Gynecology*, 50(2), 202-206.
- [17] Betebo, B., Ejajo, T., Alemseged, F., & Massa, D. (2017). Household Food Insecurity and Its Association with Nutritional Status of Children 6–59 Months of Age in East Badawacho District, South Ethiopia. *Journal of environmental and public health*, 2017.
- [18] Bhattacharya, S., Campbell, D. M., Liston, W. A., & Bhattacharya, S. (2007). Effect of body mass index on pregnancy outcomes in nulliparous women delivering singleton babies. *BMC public health*, 7(1), 168.
- [19] Bhutta, Z. A., Ahmed, T., Black, R. E., Cousens, S., Dewey, K., Giugliani, E.,... Sachdev, H. (2008). What works? Interventions for maternal and child undernutrition and survival. *The Lancet*, 371(9610), 417-440.
- [20] Bhutta, Z. A., Das, J. K., Rizvi, A., Gaffey, M. F., Walker, N., Horton, S.,... Group, T. L. N. I. R. (2013). Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *The Lancet*, 382(9890), 452-477.
- [21] Black, R. E., Allen, L. H., Bhutta, Z. A., Caulfield, L. E., de Onis, M., Ezzati, M.,... Child Undernutr, S. (2008). Maternal and child undernutrition 1 - Maternal and child undernutrition: global and regional exposures and health consequences [Review]. *Lancet*, 371(9608), 243-260. [https://doi.org/10.1016/S0140-6736\(07\)61690-0](https://doi.org/10.1016/S0140-6736(07)61690-0)
- [22] Black, R. E., Allen, L. H., Bhutta, Z. A., Caulfield, L. E., De Onis, M., Ezzati, M.,... Group, C. U. S. (2008). Maternal and child undernutrition: global and regional exposures and health consequences. *The Lancet*, 371(9608), 243-260.
- [23] Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., de Onis, M.,... Uauy, R. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries [Research Support, Non-U. S. Gov't
- [24] Review. *Lancet*, 382(9890), 427-451. [https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X)
- [25] Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., de Onis, M.,... Maternal Child Nutr Study, G. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*, 382(9890), 427-451. [https://doi.org/10.1016/S0140-6736\(13\)60937-x](https://doi.org/10.1016/S0140-6736(13)60937-x)
- [26] Black, R. E., Victora, C. G., Walker, S. P., & Maternal Child Nutr Study, G. (2013). Maternal and Child Nutrition Study Group. Maternal and child undernutrition and overweight in low-income and middle-income countries (vol 382, pg 427, 2013). *Lancet*, 382(9890), 396-396. <Go to ISI>://WOS:000322638500028
- [27] Boo, F. L., Palloni, G., & Urzua, S. (2014). Cost-benefit analysis of a micronutrient supplementation and early childhood stimulation program in Nicaragua. *Annals of the New York Academy of Sciences*, 1308(1), 139-148.
- [28] Bundara, N., Mwanri, L., & Masika, J. (2013). Addressing childhood under nutrition in Tanzania: Challenges and opportunities. *African Journal of Food, Agriculture, Nutrition and Development*, 13(1), 7288-7306.
- [29] Carletto, G., Ruel, M., Winters, P., & Zezza, A. (2015). Farm-level pathways to improved nutritional status: introduction to the special issue. *The Journal of Development Studies*.
- [30] Caulfield, L. E., de Onis, M., Blössner, M., & Black, R. E. (2004). Undernutrition as an underlying cause of child deaths associated with diarrhea, pneumonia, malaria, and measles. *The American journal of clinical nutrition*, 80(1), 193-198.
- [31] Chegere, M. J., & Stage, J. (2020). Agricultural production diversity, dietary diversity and nutritional status: Panel data evidence from Tanzania. *World Development*, 129, 104856.
- [32] Chikungu, L. C., & Madise, N. J. (2014). Seasonal variation of child under nutrition in Malawi: is seasonal food availability an important factor? Findings from a national level cross-sectional study. *Bmc Public Health*, 14, Article 1146. <https://doi.org/10.1186/1471-2458-14-1146>

- [33] Chirande, L., Charwe, D., Mbwana, H., Victor, R., Kimboka, S., Issaka, A. I., Agho, K. E. (2015). Determinants of stunting and severe stunting among under-fives in Tanzania: evidence from the 2010 cross-sectional household survey. *BMC pediatrics*, 15(1), 165.
- [34] Chuwa, M., & Mgaya, B. B. (2013). FACTORS HINDERING BREASTFEEDING PRACTICES AMONG MOTHERS IN RURAL TANZANIA. *African Journal of Midwifery & Women's Health*, 7(2).
- [35] Currie, J., & Ludwig, D. (2009). The Relationship Between Pregnancy Weight Gain and Birth Weight: A Within Family Comparison.
- [36] Darapheak, C., Takano, T., Kizuki, M., Nakamura, K., & Seino, K. (2013). Consumption of animal source foods and dietary diversity reduce stunting in children in Cambodia. *International archives of medicine*, 6(1), 1.
- [37] Dawkins, N., Shosho, N., Mamiro, P., & Pace, R. (2015). Food Insecurity and Maternal Education on Complementary Feeding Practices in a Rural Village in Tanzania. *The FASEB Journal*, 29(1 Supplement), 585.525.
- [38] De Onis, M., & Branca, F. (2016). Childhood stunting: a global perspective. *Maternal and Child Nutrition*, 12(S1), 12-26.
- [39] Dewey, K. G. (2013). The Challenge of Meeting Nutrient Needs of Infants and Young Children during the Period of Complementary Feeding: An Evolutionary Perspective [Article]. *Journal of Nutrition*, 143(12), 2050-2054. <https://doi.org/10.3945/jn.113.182527>
- [40] Disha, A. D., Rawat, R., Subandoro, A., & Menon, P. (2012). Infant and young child feeding (IYCF) practices in Ethiopia and Zambia and their association with child nutrition: Analysis of demographic and health survey data. *African Journal of Food, Agriculture, Nutrition and Development*, 12(2), 5895-5914.
- [41] Duncan, E., Ashton, L., Abdulai, A., Sawadogo-Lewis, T., King, S., Fraser, E.,... Roberton, T. (2022). Connecting the food and agriculture sector to nutrition interventions for improved health outcomes. *Food Security*, 14(3), 657-675.
- [42] Exavery, A., Kanté, A. M., Hingora, A., & Phillips, J. F. (2015). Determinants of early initiation of breastfeeding in rural Tanzania. *International breastfeeding journal*, 10(1), 27.
- [43] Faber, M., Laubscher, R., & Berti, C. (2014). Poor dietary diversity and low nutrient density of the complementary diet for 6-to 24-month-old children in urban and rural KwaZulu-Natal, South Africa. *Maternal & child nutrition*.
- [44] FAO. (2000). *The State of Food Insecurity in the World 2000: When people live with hunger and fear starvation*. FAO.
- [45] FAO. (2009, 16 -18 November). Declaration of the World Summit on Food Security. World Summit on Food Security Rome.
- [46] FAO. (2012). *International Scientific Symposium on Food and Nutrition Security information: from Valid Measurement to Effective Decision Making*.
- [47] FAO. (2016). *THE STATE OF FOOD AND AGRICULTURE: CLIMATE CHANGE, AGRICULTURE AND FOOD SECURITY*.
- [48] FAO. (2022). The state of food security and nutrition in the world 2022. Repurposing food and agricultural policies to make healthy diets more affordable.
- [49] Fawzi, W. W., Herrera, M. G., Nestel, P., Amin, A. E., & Mohamed, K. A. (1998). A longitudinal study of prolonged breastfeeding in relation to child undernutrition. *International journal of epidemiology*, 27(2), 255-260.
- [50] Ferro-Luzzi, A., Morris, S. S., Taffesse, S., Demissie, T., & D Amato, M. (2002). Seasonal undernutrition in rural Ethiopia: Magnitude, correlates, and functional significance. *FOOD AND NUTRITION BULLETIN-UNITED NATIONS UNIVERSITY*, 23(2), 227-228.
- [51] Frankenberger, T., Oshaug, A., & Smith, L. (1997). A definition of nutrition security CARE memo. *CARE, Atlanta, GA*.
- [52] Galani, Y., Orfila, C., & Gong, Y. (2022). A review of micronutrient deficiencies and analysis of maize contribution to nutrient requirements of women and children in Eastern and Southern Africa. *Critical Reviews in Food Science and Nutrition*, 62(6), 1568-1591.
- [53] Gewa, C. A., Oguttu, M., & Yandell, N. S. (2012). Maternal nutrition in rural Kenya: health and socio-demographic determinants and its association with child nutrition. *Maternal and Child Nutrition*, 8(3), 275-286. <https://doi.org/10.1111/j.1740-8709.2011.00322.x>
- [54] Ghosh, S., & Shah, D. (2004). Nutritional problems in urban slum children. *Indian pediatrics*, 41(7), 682-696.
- [55] Gillespie, S., & van den Bold, M. (2017). Agriculture, Food Systems, and Nutrition: Meeting the Challenge. *Global Challenges*.
- [56] Gol, R. M., Kheirouri, S., & Alizadeh, M. (2022). Association of dietary diversity with growth outcomes in infants and children aged under 5 years: a systematic review. *Journal of Nutrition Education and Behavior*, 54(1), 65-83.
- [57] Gomes, G. P., & Gubert, M. B. (2012). Breastfeeding in children under 2 years old and household food and nutrition security status [Article]. *Jornal De Pediatria*, 88(3), 279-282. <https://doi.org/10.2223/jped.2173>
- [58] Gould, J. F. (2017). Complementary Feeding, Micronutrients and Developmental Outcomes of Children. In *Complementary Feeding: Building the Foundations for a Healthy Life* (Vol. 87, pp. 13-28). Karger Publishers.
- [59] Haddad, L. (2013). From Nutrition Plus to Nutrition Driven: How to realize the elusive potential of agriculture for nutrition? *Food and Nutrition Bulletin*, 34(1), 39-44.
- [60] Hammond, R. A., & Dubé, L. (2012). A systems science perspective and transdisciplinary models for food and nutrition security. *Proceedings of the National Academy of Sciences*, 109(31), 12356-12363.
- [61] Han, X., Ding, S., Lu, J., & Li, Y. (2022). Global, regional, and national burdens of common micronutrient deficiencies from 1990 to 2019: A secondary trend analysis based on the Global Burden of Disease 2019 study. *EclinicalMedicine*, 44.
- [62] Hasan, M. M., Kader, A., Asif, C. A. A., & Talukder, A. (2023). Seasonal variation in the association between household food insecurity and child undernutrition in Bangladesh: Mediating role of child dietary diversity. *Maternal & child nutrition*, 19(2), e13465.

- [63] Hashim, T. H., Mgongo, M., Katanga, J., Uriyo, J. G., Damian, D. J., Stray-Pedersen, B., Msuya, S. E. (2017). Predictors of appropriate breastfeeding knowledge among pregnant women in Moshi Urban, Tanzania: a cross-sectional study. *International breastfeeding journal*, 12(1), 11.
- [64] Heidkamp, R., Ayoya, M. A., Teta, I. N., Stoltzfus, R. J., & Marhoney, J. P. (2015). Breastfeeding practices and child growth outcomes in Haiti: an analysis of data from Demographic and Health Surveys. *Maternal & child nutrition*, 11(4), 737-748.
- [65] Heidkamp, R. A., Ayoya, M. A., Teta, I. N., Stoltzfus, R. J., & Marhoney, J. P. (2015). Complementary feeding practices and child growth outcomes in Haiti: an analysis of data from Demographic and Health Surveys. *Maternal & child nutrition*, 11(4), 815-828.
- [66] Hetherington, J. B., Wiethoelter, A. K., Negin, J., & Mor, S. M. (2017). Livestock ownership, animal source foods and child nutritional outcomes in seven rural village clusters in Sub-Saharan Africa. *Agriculture & Food Security*, 6, 1-11.
- [67] Hillbruner, C., & Egan, R. (2008). Seasonality, household food security, and nutritional status in Dinajpur, Bangladesh. *Food and Nutrition Bulletin*, 29(3), 221-231.
- [68] Hirvonen, K., Taffesse, A. S., & Hassen, I. W. (2016). Seasonality and household diets in Ethiopia. *Public Health Nutrition*, 19(10), 1723-1730.
- [69] Hoddinott, J., Headey, D., & Dereje, M. (2015). Cows, missing milk markets, and nutrition in rural Ethiopia. *The Journal of Development Studies*, 51(8), 958-975.
- [70] Hotz, C., Loechl, C., de Brauw, A., Eozenou, P., Gilligan, D., Moursi, M.,... Meenakshi, J. V. (2012). A large-scale intervention to introduce orange sweet potato in rural Mozambique increases vitamin A intakes among children and women. *British Journal of Nutrition*, 108(1), 163-176.
- [71] Hotz, C., Loechl, C., Lubowa, A., Tumwine, J. K., Ndeezi, G., Masawi, A. N.,... Meenakshi, J. V. (2012). Introduction of β -carotene-rich orange sweet potato in rural Uganda resulted in increased vitamin A intakes among children and women and improved vitamin A status among children. *The Journal of nutrition*, 142(10), 1871-1880.
- [72] Huybregts, L., Houngbé, F., Salpéteur, C., Brown, R., Roberfroid, D., Ait-Aissa, M., & Kolsteren, P. (2012). The effect of adding ready-to-use supplementary food to a general food distribution on child nutritional status and morbidity: a cluster-randomized controlled trial. *PLoS medicine*, 9(9), e1001313.
- [73] IFPRI. (2016). *Global Nutrition Report 2016: From Promise to Impact: Ending Malnutrition by 2030*.
- [74] Jana, S., Frieder, G., Srijna, J., & Stefan, S. (2014). *Food Security Impact Assessment in Tanzania*.
- [75] Jones, A. D., Shrinivas, A., & Bezner-Kerr, R. (2014). Farm production diversity is associated with greater household dietary diversity in Malawi: findings from nationally representative data. *Food Policy*, 46, 1-12.
- [76] Kazaura, M. (2016). Exclusive breastfeeding practices in the Coast Region, Tanzania. *African health sciences*, 16(1), 44.
- [77] Keats, E. C., Das, J. K., Salam, R. A., Lassi, Z. S., Imdad, A., Black, R. E., & Bhutta, Z. A. (2021). Effective interventions to address maternal and child malnutrition: an update of the evidence. *The Lancet Child & Adolescent Health*, 5(5), 367-384.
- [78] Kemoe, L., Lanci, L., Mitra, P., Muehlschlegel, T., Okou, C., Spray, J.,... Unsal, F. (2022). Climate Change and Chronic Food Insecurity in Sub-Saharan Africa.
- [79] Kimani-Murage, E. W., Wekesah, F., Wanjohi, M., Kyobutungi, C., Ezech, A. C., Musoke, R. N.,... Griffiths, P. (2015). Factors affecting actualisation of the WHO breastfeeding recommendations in urban poor settings in Kenya [Article]. *Maternal and Child Nutrition*, 11(3), 314-332. <https://doi.org/10.1111/mcn.12161>
- [80] Kinabo, J. L., Mwanri, A. W., Mamiro, P. S., Kulwa, K., Bundala, N. H., Picado, J.,... Mzimhiri, R. (2017). Infant and young child feeding practices on Unguja Island in Zanzibar, Tanzania: a ProPAN based analysis. *Tanzania Journal of Health Research*, 19(3).
- [81] Kubuga, C., Lee, K., Song, S., & Song, W. O. (2016). The association between dietary diversity score and iron deficiency anemia among reproductive-aged women in Ghana. *The FASEB Journal*, 30(1 Supplement), 899.811-899.811.
- [82] Kulwa, K. B., Kinabo, J. L., & Modest, B. (2006). Constraints on good child-care practices and nutritional status in urban Dar-es-Salaam, Tanzania. *Food and Nutrition Bulletin*, 27(3), 236-244.
- [83] Kulwa, K. B., Mamiro, P. S., Kimanya, M. E., Mziray, R., & Kolsteren, P. W. (2015). Feeding practices and nutrient content of complementary meals in rural central Tanzania: implications for dietary adequacy and nutritional status. *BMC pediatrics*, 15(1), 1.
- [84] Kumar, N., Harris, J., & Rawat, R. (2015). If they grow it, will they eat and grow? Evidence from Zambia on agricultural diversity and child undernutrition. *The journal of development studies*, 51(8), 1060-1077.
- [85] Lamichhane, D. K., Leem, J. H., Kim, H. C., Park, M. S., Lee, J. Y., Moon, S. H., & Ko, J. K. (2016). Association of infant and young child feeding practices with under-nutrition: evidence from the Nepal Demographic and Health Survey. *Paediatrics and International Child Health*, 36(4), 260-269.
- [86] Lartey, A. (2015). End hunger, achieve food security and improved nutrition and promote sustainable agriculture. *UN Chronicle*, 51(4), 6-8.
- [87] Lumbanraja, S., Lutan, D., & Usman, I. (2013). Maternal weight gain and correlation with birth weight infants. *Procedia-Social and Behavioral Sciences*, 103, 647-656.
- [88] M'Kaibi, F. K., Steyn, N. P., Ochola, S. A., & Du Plessis, L. (2017). The relationship between agricultural biodiversity, dietary diversity, household food security, and stunting of children in rural Kenya. *Food science & nutrition*, 5(2), 243-254.
- [89] Mahmudiono, T., Sumarmi, S., & Rosenkranz, R. R. (2013). Household dietary diversity and child stunting in East Java, Indonesia. *Asia Pacific Journal of Clinical Nutrition*.
- [90] Maonga, A. R., Mahande, M. J., Damian, J., & Msuya, S. E. (2016). Factors Affecting Exclusive Breastfeeding among Women in Muheza District Tanga Northeastern Tanzania: A Mixed Method Community Based Study [Article]. *Maternal and Child Health Journal*, 20(1), 77-87. <https://doi.org/10.1007/s10995-015-1805-z>

- [91] Marques, A. H., O'Connor, T. G., Roth, C., Susser, E., & Bjørke-Monsen, A.-L. (2013). The influence of maternal prenatal and early childhood nutrition and maternal prenatal stress on offspring immune system development and neurodevelopmental disorders. *Frontiers in neuroscience*, 7.
- [92] Marriott, B. P., White, A., Hadden, L., Davies, J. C., & Wallingford, J. C. (2012). World Health Organization (WHO) infant and young child feeding indicators: associations with growth measures in 14 low-income countries [Article]. *Maternal and Child Nutrition*, 8(3), 354-370. <https://doi.org/10.1111/j.1740-8709.2011.00380.x>
- [93] Marriott, B. P., White, A. J., Hadden, L., Davies, J. C., & Wallingford, J. C. (2010). How well are infant and young child World Health Organization (WHO) feeding indicators associated with growth outcomes? An example from Cambodia. *Maternal & child nutrition*, 6(4), 358-373.
- [94] Marshall, N. E., Abrams, B., Barbour, L. A., Catalano, P., Christian, P., Friedman, J. E.,... Oken, E. (2022). The importance of nutrition in pregnancy and lactation: lifelong consequences. *American journal of obstetrics and gynecology*, 226(5), 607-632.
- [95] Martin, R. M. (2001). Commentary: does breastfeeding for longer cause children to be shorter? *International journal of epidemiology*, 30(3), 481-484.
- [96] Mason, J. B., Saldanha, L. S., Ramakrishnan, U., Lowe, A., Noznesky, E. A., Girard, A. W.,... Martorell, R. (2012). Opportunities for improving maternal nutrition and birth outcomes: synthesis of country experiences. *Food and Nutrition Bulletin*, 33(2_suppl1), S104-S137.
- [97] Massawe, G. D. (2017). *Farming systems and household food security in Tanzania: the case of Mvomero and Kishapu districts*. University College Dublin (Ireland).
- [98] Masset, E., Haddad, L., Cornelius, A., & Isaza-Castro, J. (2012). Effectiveness of agricultural interventions that aim to improve nutritional status of children: systematic review. *BMJ*, 344, d8222.
- [99] Masuke, R., Msuya, S. E., Mahande, J. M., Diarz, E. J., Stray-Pedersen, B., Jahanpour, O., & Mgongo, M. (2021). Effect of inappropriate complementary feeding practices on the nutritional status of children aged 6-24 months in urban Moshi, Northern Tanzania: Cohort study. *PloS one*, 16(5), e0250562.
- [100] Mavalankar, D. V., Gray, R. H., & Trivedi, C. (1992). Risk factors for preterm and term low birthweight in Ahmedabad, India. *International journal of epidemiology*, 21(2), 263-272.
- [101] Mayorga-Martínez, A. A., Kucha, C., Kwofie, E., & Ngadi, M. (2023). Designing nutrition-sensitive agriculture (NSA) interventions with multi-criteria decision analysis (MCDA): a review. *Critical Reviews in Food Science and Nutrition*, 1-20.
- [102] McLean, E., Cogswell, M., Egli, I., Wojdyla, D., & De Benoist, B. (2009). Worldwide prevalence of anaemia, WHO vitamin and mineral nutrition information system, 1993–2005. *Public Health Nutrition*, 12(04), 444-454.
- [103] Meeker, J., & Haddad, L. (2013). A state-of-the-art review of agriculture-nutrition linkages.
- [104] Menon, P., Bamezai, A., Subandoro, A., Ayoya, M. A., & Aguayo, V. (2015). Age - appropriate infant and young child feeding practices are associated with child nutrition in India: insights from nationally representative data. *Maternal & child nutrition*, 11(1), 73-87.
- [105] Metgud, C., Naik, V., & Mallapur, M. (2012). Effect of maternal nutrition on birth weight of newborn-a community-based study. *Journal of Family Welfare*, 58(2), 35-39.
- [106] Mgongo, M., Chotta, N. A., Hashim, T. H., Uriyo, J. G., Damian, D. J., Stray-Pedersen, B.,... Vangen, S. (2017). Underweight, Stunting and Wasting among Children in Kilimanjaro Region, Tanzania; a Population-Based Cross-Sectional Study. *International journal of environmental research and public health*, 14(5), 509.
- [107] Mora, J. O., & Nestel, P. S. (2000). Improving prenatal nutrition in developing countries: strategies, prospects, and challenges. *The American journal of clinical nutrition*, 71(5), 1353s-1363s.
- [108] Mrema, J. D., Elisaria, E., Mwanri, A. W., & Nyaruhucha, C. M. (2021). Prevalence and determinants of undernutrition among 6-to 59-months-old children in lowland and highland areas in Kilosa District, Tanzania: a cross-sectional study. *Journal of nutrition and metabolism*, 2021.
- [109] Muhimbula, H. S., & Issa-Zacharia, A. (2010). Persistent child malnutrition in Tanzania: Risks associated with traditional complementary foods (A review). *African Journal of Food Science*, 4(11), 679-692.
- [110] Mulu, E., & Mengistie, B. (2017). Household food insecurity and its association with nutritional status of under five children in Sekela District, Western Ethiopia: a comparative cross-sectional study. *BMC Nutrition*, 3(1), 35.
- [111] Munisi, D. Z., Buza, J., Mpolya, E. A., & Kinung'hi, S. M. (2016). Schistosoma mansoni Infections, Undernutrition and Anaemia among Primary Schoolchildren in Two Onshore Villages in Rorya District, North-Western Tanzania. *Plos One*, 11(12), e0167122.
- [112] Musumba, M., Palm, C. A., Komarek, A. M., Mutuo, P. K., & Kaya, B. (2022). Household livelihood diversification in rural Africa. *Agricultural economics*, 53(2), 246-256.
- [113] Mutisya, M., Kandala, N.-b., Ngware, M. W., & Kabiru, C. W. (2015). Household food (in) security and nutritional status of urban poor children aged 6 to 23 months in Kenya. *BMC public health*, 15(1), 1052.
- [114] NBS and ICF Macro. (2015-16). *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-MIS) 2015-16*.
- [115] Netting, M. J., & Makrides, M. (2017). Complementary Foods: Guidelines and Practices. In *Complementary Feeding: Building the Foundations for a Healthy Life* (Vol. 87, pp. 1-12). Karger Publishers.
- [116] Nisbett, N., Gillespie, S., Haddad, L., & Harris, J. (2014). Why worry about the politics of childhood undernutrition? *World Development*, 64, 420-433.
- [117] Nkonde, C., Audain, K., Kiwanuka - Lubinda, R. N., & Marinda, P. (2021). Effect of agricultural diversification on dietary diversity in rural households with children under 5 years of age in Zambia. *Food Science & Nutrition*, 9(11), 6274-6285.
- [118] Nordang, S., Shoo, T., Holmboe-Ottesen, G., Kinabo, J., & Wandel, M. (2015). Women's work in farming, child feeding practices and nutritional status among under-five children in rural Rukwa, Tanzania. *British Journal of Nutrition*, 114(10), 1594-1603. <https://doi.org/10.1017/s0007114515003116>

- [119] Nti, C. A., & Lartey, A. (2008). Influence of care practices on nutritional status of Ghanaian children. *Nutrition research and practice*, 2(2), 93-99.
- [120] Olney, D. K., Pedehombga, A., Ruel, M. T., & Dillon, A. (2015). A 2-year integrated agriculture and nutrition and health behavior change communication program targeted to women in Burkina Faso reduces anemia, wasting, and diarrhea in children 3–12.9 months of age at baseline: a cluster-randomized controlled trial. *The Journal of nutrition*, 145(6), 1317-1324.
- [121] Osei, A., Pandey, P., Spiro, D., Nielson, J., Shrestha, R., Talukder, Z., Haselow, N. (2010). Household food insecurity and nutritional status of children aged 6 to 23 months in Kailali District of Nepal. *Food and Nutrition Bulletin*, 31(4), 483-494.
- [122] Pobee, R., Akonor, P., & Bonsi, E. (2017). Orange-fleshed sweet potato based complementary food provides sufficient vitamin A for infants aged 6-12 months.
- [123] Psaki, S., Bhutta, Z. A., Ahmed, T., Ahmed, S., Bessong, P., Islam, M.,... Nesamvuni, C. (2012). Household food access and child malnutrition: results from the eight-country MAL-ED study. *Population health metrics*, 10(1), 24.
- [124] Rah, J., Akhter, N., Semba, R., De Pee, S., Bloem, M., Campbell, A.,... Kraemer, K. (2010). Low dietary diversity is a predictor of child stunting in rural Bangladesh. *European Journal of Clinical Nutrition*, 64(12), 1393-1398.
- [125] Ramakrishnan, U., Young, M. F., & Martorell, R. (2017). Maternal Nutrition and Birth Outcomes. In *Nutrition and Health in a Developing World* (pp. 487-502). Springer.
- [126] Ringler, C., Zhu, T., Cai, X., Koo, J., & Wang, D. (2010). Climate change impacts on food security in sub-Saharan Africa. *Insights from Comprehensive Climate Change Scenarios*.
- [127] Risnes, K. R., Vatten, L. J., Baker, J. L., Jameson, K., Sovio, U., Kajantie, E.,... Painter, R. C. (2011). Birthweight and mortality in adulthood: a systematic review and meta-analysis. *International journal of epidemiology*, 40(3), 647-661.
- [128] Ritchie, H., & Roser, M. (2017). Micronutrient deficiency. *Our World in data*.
- [129] Roba, K. T., O'Connor, T. P., Belachew, T., & O'Brien, N. M. (2016). Variations between post-and pre-harvest seasons in stunting, wasting, and Infant and Young Child Feeding (IYCF) practices among children 6-23 months of age in lowland and midland agro-ecological zones of rural Ethiopia. *The Pan African medical journal*, 24.
- [130] Ruel, M. T. (2002). *Is dietary diversity an indicator of food security or dietary quality?*
- [131] Ruel, M. T., Alderman, H., Maternal, & Group, C. N. S. (2013). Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *The Lancet*, 382(9891), 536-551.
- [132] Ruel, M. T., Quisumbing, A. R., & Balagamwala, M. (2018). Nutrition-sensitive agriculture: what have we learned so far? *Global food security*, 17, 128-153.
- [133] Saaka, M., & Osman, S. M. (2013). Does household food insecurity affect the nutritional status of preschool children aged 6–36 Months? *International Journal of Population Research*, 2013.
- [134] Saaka, M., & Rauf, A. A. (2015). Role of dietary diversity in ensuring adequate haematological status during pregnancy. *International Journal of Medical Research & Health Sciences*, 4(4), 749-755.
- [135] Saaka, M., Wemakor, A., Abizari, A.-R., & Aryee, P. (2015). How well do WHO complementary feeding indicators relate to nutritional status of children aged 6–23 months in rural Northern Ghana? *BMC public health*, 15(1), 1.
- [136] Saha, K. K., Frongillo, E. A., Alam, D. S., Arifeen, S. E., Persson, L. Å., & Rasmussen, K. M. (2009). Household food security is associated with growth of infants and young children in rural Bangladesh. *Public Health Nutrition*, 12(09), 1556-1562.
- [137] Santos, L. P. d., & Gigante, D. P. (2013). Relationship between food insecurity and nutritional status of Brazilian children under the age of five. *Revista Brasileira de Epidemiologia*, 16(4), 984-994.
- [138] Scaling Up Nutrition. (2012). *UNITED NATIONS SYSTEMS. STANDING COMMITTEE ON NUTRITION. 2012. 'Scaling up Nutrition Strategy 2012-2015'*
- [139] Scrimshaw, N. S. (2010). INCAP studies of nutrition and infection. *Food and Nutrition Bulletin*, 31(1), 54-67.
- [140] Semali, I. A., Tengia-Kessy, A., Mmbaga, E. J., & Leyna, G. (2015). Prevalence and determinants of stunting in under-five children in central Tanzania: remaining threats to achieving Millennium Development Goal 4. *BMC public health*, 15(1), 1153.
- [141] Sheffield, P. E., & Landrigan, P. J. (2011). Global climate change and children's health: threats and strategies for prevention. *Environmental health perspectives*, 119(3), 291.
- [142] Shetty, P. (2015). From food security to food and nutrition security: role of agriculture and farming systems for nutrition. *Current Science*, 109, 456-461.
- [143] Shirima, C. P., Kimanya, M. E., Routledge, M. N., Srey, C., Kinabo, J. L., Humpf, H.-U.,... Gong, Y. Y. (2015). A prospective study of growth and biomarkers of exposure to aflatoxin and fumonisin during early childhood in Tanzania. *Environmental health perspectives*, 123(2), 173.
- [144] Shrimpton, R., Victora, C. G., de Onis, M., Lima, R. C., Blössner, M., & Clugston, G. (2001). Worldwide timing of growth faltering: implications for nutritional interventions. *Pediatrics*, 107(5), e75-e75.
- [145] Singh, A., Singh, A., & Ram, F. (2014). Household food insecurity and nutritional status of children and women in Nepal. *Food and Nutrition Bulletin*, 35(1), 3-11.
- [146] Slavchevska, V. (2015). Agricultural production and the nutritional status of family members in Tanzania. *The Journal of Development Studies*, 51(8), 1016-1033.
- [147] Smith, L. C., & Haddad, L. (2015). Reducing child undernutrition: past drivers and priorities for the post-MDG era. *World Development*, 68, 180-204.
- [148] Stevens, G. A., Beal, T., Mbuya, M. N., Luo, H., Neufeld, L. M., Addo, O. Y., Brown, K. H. (2022). Micronutrient deficiencies among preschool-aged children and women of reproductive age worldwide: a pooled analysis of individual-level data from population-representative surveys. *The Lancet Global Health*, 10(11), e1590-e1599.

- [149] Stevens, G. A., Finucane, M. M., De-Regil, L. M., Paciorek, C. J., Flaxman, S. R., Branca, F., Nutr Impact Model Study, G. (2013). Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population-representative data [Article]. *Lancet Global Health*, 1(1), E16-E25. <Go to ISI>://WOS:000336422200009
- [150] Stewart, C. P., Iannotti, L., Dewey, K. G., Michaelsen, K. F., & Onyango, A. W. (2013). Contextualising complementary feeding in a broader framework for stunting prevention [Article]. *Maternal and Child Nutrition*, 9, 27-45. <https://doi.org/10.1111/mcn.12088>
- [151] Suseela, R., Behrman, J. A., Menon, P., & Quisumbing, A. R. (2012). Gender: A Key Dimension Linking Agricultural Programs to Improved Nutrition and Health.
- [152] Swaminathan, M. (2014). Zero hunger. *Science*, 345(6196), 491-491.
- [153] Tabrizi, F. M., & Barjasteh, S. (2015). Maternal hemoglobin levels during pregnancy and their association with birth weight of neonates. *Iranian journal of pediatric hematology and oncology*, 5(4), 211.
- [154] Tafese, Z., & Kebebu, A. (2017). A systematic review of maternal feeding practice and its outcome in developing countries.
- [155] Terefe, B., Birhanu, A., Nigussie, P., & Tsegaye, A. (2015). Effect of maternal iron deficiency anemia on the iron store of newborns in Ethiopia. *Anemia*, 2015.
- [156] Ubeysekara, N. H., Jayathissa, R., & Wijesinghe, C. J. (2015). Nutritional status and associated feeding practices among children aged 6-24 months in a selected community in Sri Lanka: a cross sectional study. *Eur J Prev Med*, 3(2-1), 15-23.
- [157] Udoh, E. E., & Amodu, O. K. (2016). Complementary feeding practices among mothers and nutritional status of infants in Akpabuyo Area, Cross River State Nigeria. *SpringerPlus*, 5(1), 2073.
- [158] UNICEF. (1998). *The state of the world's children 1998: Focus on nutrition*. Oxford University Press.
- [159] UNICEF. (2013). *Improving child nutrition: the achievable imperative for global progress*.
- [160] UNICEF. (2015a). *The State of the World's Children 2015: Reimagine the Future: Innovation for Every Child*.
- [161] UNICEF. (2015b). *UNICEF's approach to scaling up nutrition for mothers and their children. Discussion paper* (Programme Division, Issue).
- [162] UNICEF. (2016). *FROM THE FIRST HOUR OF LIFE: Making the case for improved infant and young child feeding everywhere*. UNICEF.
- [163] UNICEF, W. a. (2003). *The global strategy on infant and young child nutrition (IYCN)* W. H. Organization.
- [164] UNICEF, W. W. B. (2023). Levels and trends in child malnutrition: UNICEF/WHO/World Bank Group joint child malnutrition estimates: key findings of the 2023 edition. In *Levels and trends in child malnutrition: UNICEF/WHO/World Bank Group joint child malnutrition estimates: key findings of the 2023 edition*. UNICEF, New York; WHO, Geneva; World Bank, Washington, <https://www.who.int/publications/i/item/9789240073791>
- [165] Victor, R., Baines, S. K., Agho, K. E., & Dibley, M. J. (2014). Factors associated with inappropriate complementary feeding practices among children aged 6-23 months in Tanzania. *Maternal and Child Nutrition*, 10(4), 545-561. <https://doi.org/10.1111/j.1740-8709.2012.00435.x>
- [166] Victora, C. G., Bahl, R., Barros, A. J., França, G. V., Horton, S., Krasevec, J.,... Rollins, N. C. (2016). Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *The Lancet*, 387(10017), 475-490.
- [167] Victora, C. G., Bahl, R., Barros, A. J. D., Franca, G. V. A., Horton, S., Krasevec, J.,... Lancet Breastfeeding Series, G. (2016). Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect [Article]. *Lancet*, 387(10017), 475-490. <Go to ISI>://WOS:000368986800036
- [168] Victora, C. G., Christian, P., Vidaletti, L. P., Gatica-Domínguez, G., Menon, P., & Black, R. E. (2021). Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda. *The Lancet*, 397(10282), 1388-1399.
- [169] Victora, C. G., de Onis, M., Hallal, P. C., Blössner, M., & Shrimpton, R. (2010). Worldwide timing of growth faltering: revisiting implications for interventions. *Pediatrics*, 125(5), 1519-1519.
- [170] Von Grebmer, K., Bernstein, J., Resnick, D., Wiemers, M., Reiner, L., Bachmeier, M.,... Foley, C. (2022). Global hunger index: Food systems transformation and local governance. *Welt Hunger Hilfe CONCERNWorldwide: Bonn/Dublin, Ireland*.
- [171] Wagah, M., Hodge, J., & Lewis, A. (2015). Leveraging Agriculture for Nutrition in East Africa (LANEA). In.
- [172] Webb - Girard, A., Cherobon, A., Mbugua, S., Kamau - Mbuthia, E., Amin, A., & Sellen, D. W. (2012). Food insecurity is associated with attitudes towards exclusive breastfeeding among women in urban Kenya. *Maternal & child nutrition*, 8(2), 199-214.
- [173] Weingärtner, L. (2009). The concept of food and nutrition security. *Achieving food and nutrition security*, 3, 21-52.
- [174] Wenban-Smith, H. (2014). *Rural-Urban Linkages: Tanzania Case Study*.
- [175] WHO. (1995). *Physical status: The use of and interpretation of anthropometry, Report of a WHO Expert Committee* (WHO technical report series no 854, Issue).
- [176] WHO. (2008). *Indicators for assessing infant and young child feeding practices, part 1: definitions; conclusions of a consensus meeting held 6-8 November 2007*.
- [177] WHO. (2010). *Indicators for Assessing Infant and Young Child Feeding Practices. Part II: Measurement*.
- [178] WHO. (2021). Indicators for assessing infant and young child feeding practices: definitions and measurement methods.
- [179] Wijesinha-Bettoni, R., Kennedy, G., Dirorimwe, C., & Muehlhoff, E. (2013). Considering seasonal variations in food availability and caring capacity when planning complementary feeding interventions in developing countries. *International Journal of Child Health and Nutrition*, 2(4), 335-352.

- [180] Wordofa, M. G., & Sassi, M. (2020). Impact of agricultural interventions on food and nutrition security in Ethiopia: uncovering pathways linking agriculture to improved nutrition. *Cogent food & agriculture*, 6(1), 1724386.
- [181] Woodhill, J., Kishore, A., Njuki, J., Jones, K., & Hasnain, S. (2022). Food systems and rural wellbeing: challenges and opportunities. *Food Security*, 14(5), 1099-1121.
- [182] Young, S. L., Plenty, A. H., Luwedde, F. A., Natamba, B. K., Natureeba, P., Achan, J., Osterbauer, B. (2014). Household food insecurity, maternal nutritional status, and infant feeding practices among HIV-infected Ugandan women receiving combination antiretroviral therapy. *Maternal and Child Health Journal*, 18(9), 2044-2053.
- [183] Zongrone, A., Winskell, K., & Menon, P. (2012). Infant and young child feeding practices and child undernutrition in Bangladesh: insights from nationally representative data. *Public Health Nutrition*, 15(9), 1697-1704.